arXiv:1402.0135v1 [math.OA] 1 Feb 2014

Property RD and the Classification of Traces on Reduced Group C^*-algebras of Hyperbolic Groups

Sherry Gong

February 4, 2014

Abstract

In this paper, we show that if G by a non-elementary word hyperbolic group and $a \in G$ an element, if the conjugacy class of a is infinite, then all traces $\tau : C^*_{\text{red}}(G) \to \mathbb{C}$ vanish on a. We show that all traces $\theta : C^*_{\text{red}}(G) \to \mathbb{C}$ are linear combinations of traces $\chi_g : C^*_{\text{red}}(G) \to \mathbb{C}$ given by

$$\chi_g = \begin{cases} 1 & g \in C(g) \\ 0 & \text{else} \end{cases}$$

where $C(g)$ is the conjugacy class of g. To do this, we introduce a new method to study traces, using Sobolev norms and property RD.

1 Introduction

Let G be a countable discrete group and let $\ell^2(G)$ denote the Hilbert space of square-summable functions $G \to \mathbb{C}$. Let $B(\ell^2(G))$ denote the algebra of bounded linear operators on $\ell^2(G)$. Note that elements of G act on $\ell^2(G)$ as bounded linear operators, so there is a natural map $\mathbb{C}[G] \to B(\ell^2(G))$, where $\mathbb{C}[G]$ denotes the group algebra of G. We define the reduced group C^*-algebra, denoted $C^*_{\text{red}}G$, to be the operator norm closure of $\mathbb{C}[G]$ in $B(\ell^2(G))$. The reduced group C^* algebra plays an important role in noncommutative geometry, and these algebras are studied in detail in the book [3].

A trace on $C^*_{\text{red}}G$ is a bounded linear map $\tau : C^*_{\text{red}}G \to \mathbb{C}$ that satisfies $\tau(xy) = \tau(yx)$ for $x, y \in C^*_{\text{red}}G$. It is easy to see that such a map satisfies that for $g, h \in G \in C^*_{\text{red}}G$, we have $\tau(ghg^{-1}) = \tau(h)$, and also that if we had a bounded linear map $\tau : C^*_{\text{red}}G \to \mathbb{C}$ $\tau(ghg^{-1}) = \tau(h)$, then it is a trace.

In [11], Weinberger and Yu studied the degree of non-rigidity of manifolds using idempotents in C^* algebras of groups associated to torsion elements in those groups, and exploring whether they are linearly independent in K-theory. One approach to this is to study whether idempotents can be shown to be linearly independent using traces, as explored in [6]. The study of idempotents also leads to results in the study of the moduli space of positive scalar curvature metrics, as in [12] and [13].

The question of studying the traces on reduced group C^* algebras was introduced by Powers in [10], who proved that up to scalar multiplication, there is only one trace on $C^*_{\text{red}}G$ for non-abelian free groups G. It was also studied extensivesly in [2], [4], and [5], where it was shown that if G is a torsion-free non-elementary hyperblic group, then $C^*_{\text{red}}G$ has a only one trace, up to scalar. Similar questions were also studied in [1] for relatively hyperbolic groups.

In this paper, we introduce a new method of studying traces using property RD. Recall that a length function on a group G is a function $l : G \to \mathbb{Z}_{\geq 0}$, such that $l(fg) \leq l(f)+l(g)$, $l(g^{-1}) = l(g)$, and $l(e) = 0$, such that for $S \subset \mathbb{Z}_{\geq 0}$ finite $l^{-1}(S)$ is also finite. Given such a length function on a group, there is a norm $\| \cdot \|_{H^s}$ on $\mathbb{C}G$ given by

$$\| \sum c_g g\|^2_{H^s} = \sum |c_g|^2 (1 + l(g))^{2s}.$$

Let $H^s(G)$ denote the completion of $\mathbb{C}G$ with respect to this norm. Then we say G has rapid decay property, or property RD if for some s, there is a length function on it and a constant C such that

$$\| \sum c_g g\|_{\text{red}} \leq C \| \sum c_g g\|_{H^s}.$$

This property was introduced by Connes and studied extensively by Jolissaint in [7] and [8].

For a countable discrete group G with length function l and a subset $X \subset G$, we say X has polynomial growth if for $X_l = \{g | g \in X, l(g) = l\}$ and $n_l = |X_l|$, we have $n_l < P(l)$ for some polynomial P. If such a P does not exist, we say that X grows super-polynomially. We say X grows at least exponentially if there are a, b, c with $a, b > 0$ such that $n_l > ae^{bl}+c$.

In [6], I showed that for groups with property RD, if the conjugacy class of an element $g \in G$ grows super-polynomially, then any trace $\tau : C^*_{\text{red}}G \to \mathbb{C}$ vanishes on $g \in G$. We will explain this result in more detail in section 3.

We will use this method to classify all traces on all non-elementary hyperbolic groups, extending de la Harpe's result on torsion-free non-elementary hyperbolic groups.

The author would like to thank Prof. Guoliang Yu for his helpful advice and guidance,

and Prof. David Kerr for his suggestion to look into questions regarding unique traces on reduced group C^* algebras.

2 On conjugacy classes

In this section, we show that if a conjugacy class in a hyperbolic group is infinite, then it grows exponentially. In the next section, we will use this to find all the traces.

Let G be a finitely generate discrete group and $\delta > 0$ a real number. Consider the metric space obtained by considering the Cayley graph of G, and identifying each edge with the unit interval. Then, we say G is δ-hyperbolic if for any $x, y, z \in G$,

$$[x, y] \subset B_\delta([y, z] \cup [z, x])$$

where $[x, y]$ denotes the geodesic between x and y in the metric space, and for a subset $X \subset G$, $B_\delta(X)$ denotes points a such that there is $b \in X$ with $d(a, b) < \delta$.

We say that a group G is elementary if G contains $\mathbb{Z}$ as a finite index subgroup. We study the problem of conjugacy class growth in non-elementary hyperbolic groups.

Theorem 2.1. Let G be a non-elementary hyperbolic group and let $a \in G$ be an element. Suppose that $C(a)$, the conjugacy class of a in G is infinite. Then $C(a)$ grows at least exponentially.

Notation: For t a non-negative real number, we let B_t denote the ball in the Cayley graph of G around the identity of radius t.

Proof. To show this theorem, we want to consider some element in $x \in C(a)$ of length n, where n is large, and consider conjugating it to gxg^{-1} where $\ell(g) < \frac{n}{7}$. The number of g with $\ell(g) < \frac{n}{7}$ grows exponentially in n, and $\ell(gxg^{-1})$ is then in the range $\left[\frac{5n}{7}, \frac{9n}{7}\right]$ so if all different g in $\ell(g) < \frac{n}{7}$ gave rise to different gxg^{-1}, we would be done, because we would have shown that the conjugacy class of a has exponentially many elements in the range.

However, this is not quite true, because there could be g and h with $\ell(g), \ell(h) < \frac{n}{7}$, but so that $gxg^{-1} = hxh^{-1}$. However, we will show that this cannot happen too much. More precisely, we prove the following lemma:

Lemma 2.2. Let G be a non-elementary hyperbolic group and $a \in G$ be a fixed element. Let x be in $C(a)$, and let $n = \ell(x)$. Let $y \in C(a)$ have $\frac{5n}{7} < \ell(y) < \frac{9n}{7}$. Let $m = \ell(y)$. Let $k < \frac{n}{7}$ be an integer. Then the number of h in G such that

1. $\ell(h) = k$

2. $hxh^{-1} = y$

is $O(k)$, meaning it is bounded above by $Ck + C_1$ where C and C_1 are constants that depend only on G and a, and independent of n and k.

Thus, the number of elements $h \in G$ such that

1. $\ell(h) < \frac{n}{7}$

2. $hxh^{-1} = y$

is $O(n^2)$, meaning it is bounded by $C_2n^2 + C_3$.

Proof of Lemma. In triangle ABC in a graph, let a, b, c, denote the sides; that is, $c = \ell(AB)$, $b = \ell(AC)$ and $a = \ell(BC)$. We say that triangle ABC is δ_1-thin if for points $D \in BC$, $E \in AC$ and $F \in AB$ with $AE = AF = \frac{b+c-a}{2}$, $\ell(BD) = \ell(BF) = \frac{a+c-b}{2}$ and $\ell(CD) = \ell(CE) = \frac{a+b-c}{2}$, we have $\ell(DE), \ell(DF), \ell(EF) < \delta_1$. Proposition 2.12 in [9] says that for a δ-hyperbolic graph, there is δ_1 such that all triangles are δ_1-thin.

Let us replace δ with the max of δ and δ_1 so that now the graph is δ-hyperbolic and all triangles are δ-thin.

Let us assume $n > 5\delta$, which we may, because there are finitely many cases for $n \leq 5\delta$, so we can shove those cases into the constant C_1.



Figure 1:

Consider the geodesic picture shown in the diagram, where $A = e \in G$, $B = h^{-1}$, $C = xh^{-1}$ and $D = hxh^{-1}$, and AB, BC, CD, and AD are geodesics. Thus, BC is composed of the same sequence of $s_i \in S$ edges as the geodesic for x would be, and CD the same as the geodesic for h.

(In my picture the Cayley graph is formed by taking generators S and joining g to sg for $s \in S$. Thus, my picture is invariant under the right action of G.)

Now consider triangle ABC. Applying that it is δ-thin, we have $E \in AB$, $F \in BC$ and $G \in AC$ such that $\ell(EF), \ell(EG), \ell(FG) < \delta$, and $\ell(BE) = \ell(BF)$ and $\ell(AE) = \ell(AG)$. Now, let us apply hyperbolicity to triange ACD. We have $G \in AC$ so there is $H \in AD \cup CD$ such that $\ell(GH) < \delta$.

We claim that $H \in AD$. If not, $H \in CD$, but then $\ell(AC) \leq \ell(AG) + \ell(GH) + \ell(HC)$ and $\ell(AG) = \ell(AE) < k$, and $\ell(HC) < k$, so $\ell(AC) < 2k + \delta$, but $\ell(AC) > n - k$, so $n - k < 2k + \delta$, so $n < 3k + \delta < \frac{3n}{7} + \delta$, which is impossible because $n > 5\delta$.

Thus $H \in AD$. So we have $H \in AD$ with $\ell(EH) < \ell(EG) + \ell(GH) < 2\delta$. Also, $\ell(AE) = \ell(AG)$, so

$$\ell(AE) - \delta < \ell(AE) - \ell(GH) = \ell(AG) - \ell(GH) <$$

$$\ell(AH) < \ell(AG) + \ell(GH) = \ell(AE) + < \ell(GH) < \ell(AE) + \delta.$$

So now for fixed x and y, and for $i = \ell(AE)$, $0 \leq i \leq k$, we have fixed $F \in BC$, because F has $\ell(BF) = \ell(BE) = k - \ell(AE) = k - i$, and BC is just the h^{-1} right translate of the geodesic for x, so BF is fixed, because it is just the product of the first $k - i$ edges of the geodesic of x.

Now, BE, as a group element, has distance at most δ from BF, so there are $|B_\delta|$ possibilities for BE.

For i fixed, as above, there are 2δ possibilities for AH, because H is along AD, which is the geodesic of y, and we showed above that $i - \delta = \ell(AE) - \delta < \ell(AH) < \ell(AE) + \delta = i + \delta$. So AH is the product of the first t edges of the geodesic of y.

For each of the 2δ choices of AH, there are at most $B_{2\delta}$ choices of AE, because AE and AH are within 2δ of each other as group elements. Thus, in total, for fixed i, there are $2\delta B_{2\delta}$ possibilities for AE.

So we have, for fixed $i = \ell(AE) = k - \ell(BE)$, there are B_δ choices of AE and $2\delta B_{2\delta}$ choices of BE so there are $2\delta B_\delta B_{2\delta}$ choices of $AB = EB \cdot AE$. But $AB = h^{-1}$, so there are at most $O(k)$ choices of h for a given k and i.

There are k choices of $i \in [0, k]$, so in total there are at most $k(2\delta B_\delta B_{2\delta})$ possible h for

a given k, giving us the first statement in the lemma.

Adding up the values for different $k < n/7$ is adding up $n/7$ summands, each of which is $O(n)$, so we get that there are $O(n^2)$ possible h with $\ell(h) < n/7$ and $hxh^{-1} = y$.

□

Now we complete the proof of the theorem. We are assuming that the conjugacy class of a is infinite, so there are $x \in C(a)$ of arbitrarily large length. For large n, consider some $x \in C(a)$ with $\ell(x) = n$. Then for $g \in B_{n/7}$, we may consider $gxg^{-1} \in C(a)$. Then $5n/7 < \ell(gxg^{-1} < 9n/7$, so we can map $B_{n/7}$ to the elements of the conjugacy class of a of length in the range $\left[\frac{5n}{7}, \frac{9n}{7}\right]$.

$$\phi : B_{n/7} \to C(a) \cap (B_{9n/7} - B_{5n/7})$$

By the above lemma, the pre-image of any element under ϕ has size at most $C_2 n^2 + C_3$, for constants C_2, C_3 depending only on a and G. But $|B_{n/7}|$ grows exponentially by a theorem in Koubi (also attached), that says that non-elementary hyperbolic groups have exponential growth. So the number of elements in the image of ϕ, grows at least with $e^{\alpha n}/n^2$, which is still exponential.

Thus the conjugacy class of $C(a) \cap B_{9n/7}$ grows exponentially in n, so $C(a)$ has exponential growth, as desired.

□

Corollary 2.3. For G a non-elementary hyperbolic group and $a \in G$ a non-torsion element, the conjugacy class of a grows exponentially.

Proof. By the theorem, it suffices to show that the conjugacy class is infinite. Let $C(a)$ be the conjugacy class and $Z(a)$ be the centraliser. Then $C(a) = G/Z(a)$, so if the conjugacy class is finite, then $Z(a) \subset G$ has finite index. But, by theorem 3.35, $Z(a)$ is a finite extension of $\langle a \rangle$, so it is virtually cyclic. But any group is quasi-isometric to any finite index subgroup, so that would mean that G is also virtually cyclic, which is impossible. □

3 On traces

In this section, we show the main result; namely, we classify all traces on non-elementary word-hyperbolic groups.

Theorem 3.1. Let G be a non-elementary word-hyperbolic group. Then the traces θ :

$C^*_{\text{red}}(G) \to \mathbb{C}$ are exactly the linear combinations of $\chi_g : C^*_{\text{red}}(G) \to \mathbb{C}$ given by

$$\chi_g = \begin{cases} 1 & g \in C(g) \\ 0 & \text{else} \end{cases}$$

where g is an element of finite conjugacy class and $C(g)$ is its conjugacy class.

Proof of Theorem 3.1. As was shown in [9] [Theorem 3.27], in a hyperbolic group, there are only finitely many conjugacy classes of torsion elements.

We will show that all infinite order elements have infinite conjugacy class. Suppose, for the sake of contradiction that there is $g \in G$ an element of infinite order with finite conjugacy class $C(g)$. Then $C(g) = G/Z(g)$, where $Z(g)$ is the centraliser of G. However, it was shown in [9] [Theorem 3.35] that $Z(g)$ is a finite extension of the subgroup generated by g, so that it is elementary. But then if $C(g)$ is finite, then G would also be elementary, a contradiction.

So we see that there are only finitely many elements that have finite conjugacy class. For each such element g, it is easy to see that the map

$$\chi_g = \begin{cases} 1 & g \in C(g) \\ 0 & \text{else} \end{cases}$$

defines a bounded linear functional $C^*_{\text{red}}(G) \to \mathbb{C}$.

Any linear combination of these finitely many traces is also a bounded trace. To do this, we will use the fact that word hyperbolic groups have property RD:

Definition 3.2. Given such a length function on a group, there is a norm $\| \cdot \|_{H^s}$ on $\mathbb{C}G$ given by

$$\| \sum c_g g \|^2_{H^s} = \sum |c_g|^2 (1 + l(g))^{2s}.$$

Let $H^s(G)$ denote the completion of $\mathbb{C}G$ with respect to this norm. Then we say G has rapid decay property, or property RD if for some s, there is a length function on it and a constant C such that

$$\| \sum c_g g \|_{\text{red}} \leq C \| \sum c_g g \|_{H^s}.$$

It was shown in [7] that all hyperbolic groups have property RD.

Now, we will apply Lemma 2.8 in [6], which states the following:

Theorem 3.3. Let G be a group with property RD and $g \in G$ be an element whose conjugacy class grows faster than polynomially. Then all traces $C^*_{\text{red}}G \to \mathbb{C}$ vanish on g.

In our case, word-hyperbolic groups have property RD, so applying Theorem 2.1, we have that all traces $C^*_{\text{red}}G \to \mathbb{C}$ vanish on all elements with infinite conjugacy class, and they must be linear combinations of the above traces, as desired.

□

In particular, this gives another proof of de la Harpe's result regarding traces on torsion-free non-elementary word-hyperbolic groups, which can be found in [4]:

Corollary 3.4. Let G be a torsion-free non-elementary word-hyperbolic group. Then there is only one trace $\tau : C^*_{\text{red}}G \to \mathbb{C}$, up to scalar multiples.

Theorem 3.1 also implies that every hyperbolic group can be written as an extension of a hyperbolic group with unique trace by a finite group, namely:

Theorem 3.5. Let G be a hyperbolic group. Then it has a finite normal subgroup N such that G/N is a hyperbolic group with unique trace, up to scalar multiples.

Proof. Let $N \subset G$ be the subset of elements with finite conjugacy class. Note that all elements in it have finite order, as we showed in the proof of Theorem 3.1.

Lemma 3.6. We claim that N is a subgroup.

Proof. It is easy to see that N is closed under inversion, so we want that if $g_1, g_2 \in H$ then $g_1 g_2 \in N$. For an element $g \in G$, let $Z(g) \subset G$ be the subgroup consisting of elements that commute with g. Then, note that $g \in N$ if and only if $Z(g)$ has finite index as a subgroup of G. Thus, it suffices to show that if $Z(g_1)$ and $Z(g_2)$ have finite index, then $Z(g_1 g_2)$ does also.

Note that $Z(g_1 g_2) \supset Z(g_1) \cap Z(g_2)$, so it suffices to show that if $H_1, H_2 \subset G$ are finite index subgroups, then $H_1 \cap H_2$ is also. There is a natural injection $H_1/H_1 \cap H_2 \to G/H_2$. Thus, $|H_1/H_1 \cap H_2| < \infty$. Let

$$G = a_1 H_1 \cup a_2 H_1 \cup \cdots \cup a_n H_1$$

and

$$H_1 = b_1(H_1 \cap H_2) \cup b_2(H_1 \cap H_2) \cup \cdots \cup b_m(H_1 \cap H_2).$$

Then $G = \bigcup_{i,j} a_i b_j (H_1 \cap H_2)$, so $H_1 \cap H_2$ has finite index, as desired. □

By the lemma, we have that N is a subgroup of G. As we showed in the proof of Theorem 3.1, N is finite. Also, it is easy to see that N is closed under conjugation. Thus, N is a finite normal subgroup.

Note that G/N is hyperbolic because hyperbolicity is preserved under quasi-isometries, and it is easy to see that the map $G \to G/N$ is a quasi isometry. Thus, G/N is hyperbolic. It remains to show that G/N has unique trace $\tau : C^*_{\text{red}}(G/N) \to \mathbb{C}$. Such a trace τ induces a trace $C^*_{\text{red}}(G) \to \mathbb{C}$, because N is a finite normal subgroup. By Theorem 3.1, the trace $C^*_{\text{red}}(G) \to \mathbb{C}$ vanishes on all elements outside of N, so the map $\tau : C^*_{\text{red}}(G/N) \to \mathbb{C}$ vanishes on all elements except the identity, as desired.

□

4 Examples

In this section, we give some examples of hyperbolic groups with torsion that have unique trace by Theorem 3.1.

Example. Let $G = G_1 * G_2$ be the free product of two non-trivial hyperbolic groups G_1 and G_2. Suppose that $|G_1| > 2$. This assumption does not lose any generality, because $\mathbb{Z}/2 * \mathbb{Z}/2$ is elementary. Then G is hyperbolic and all its conjugacy classes are infinite. It is hyperbolic because the free product of two hyperbolic groups is hyperbolic. To see that the conjugacy classes have infinite order: note that every element $x \in G$ can be uniquely expressed as $x = g_1h_1g_2h_2 \cdots g_nh_n$ where $g_i \in G_1$, $h_i \in G_2$ and $g_i \neq 1$ for $i > 1$ and $h_i \neq 1$ for $i < n$.

In the case $g_1 = 1$ and $h_n \neq 1$, then consider some element $g \neq 1 \in G_1$ and $h \neq 1 \in G_2$, and consider $hghg \cdots hghgxg^{-1}h^{-1}g^{-1}h^{-1} \cdots g^{-1}h^{-1} = (hg)^t x (hg)^{-t}$, that is conjugation of x by $(hg)^t$. As t varies amongst positive integers, $(hg)^t x (hg)^{-t}$ goes through different elements of G, all in the conjugacy class of x.

The case where $g_1 \neq 1$ and $h_n = 1$ is similar.

For the case $g_1 \neq 1$, and $h_n \neq 1$, conjugate by $(hg)^t$ where $g \in G_1$ has $g \neq 1$ and $g \neq g_1^{-1}$. Then the same argument as the above applies. Similarly, for the case $g_1 = 1$ and $h_n = 1$, pick $g \in G_1$, with $g \neq 1$ and $g \neq g_n^{-1} \in G_1$.

Thus, we have that G is a hyperbolic group with infinite conjugacy classes, so, applying Theorem 3.1, whenever such a group is non-elementary, it has unique traces.

This example shows that many hyperbolic groups with torsion elements still have unique traces, extending de la Harpe's results in [2], [4], and [5] that torsion-free non-elementary hyperbolic groups have unique traces.

Now let us give an example of a non-elementary hyperbolic group that does not have unique trace (up to scalar multiples). For this, by Theorem 3.1, it suffices that there are non-identity elements with finite conjugacy class.

There are obvious examples of such groups:

Example. Let G by a non-elementary hyperbolic group and G_1 be a non-trivial finite group. Then in $G_1 \times G$, the direct product of G_1 and G, is hyperbolic, because it is quasi-isometric to G. The elements in $G_1 \times e \subset G_1 \times G$, where e is the identity in G, have finite conjugacy class.

Here is an example of a non-elementary hyperbolic group that does not have only one trace up to scalar, but which is not the direct product of a finite group and a hyperbolic group.

Example. Let $G = \langle x, y, a \rangle / \langle a^3, aya^{-1}y^{-1}, xax^{-1}a^{-2} \rangle$. This group G is hyperbolic, because it is quasi-isometric to the free group on two generators, $\langle x, y, \rangle \subset G$. In G, a and a^2 clearly have finite conjugacy class, as their conjugacy class is $\{a, a^2\}$. Thus, by Theorem 3.1, G does not have unique trace.

However, it is not the direct product of a finite group and a hyperbolic group whose reduced C^* algebra has unique trace up to scalar multiples, because the only elements of finite order in G are a and a^2, so if G were such a direct product, it would be $\mathbb{Z}/3\mathbb{Z} \times G_1$ for some hyperbolic group G_1. But in that case, a would not be conjugate to a^2.

References

[1] G.N. Arzhantseva, A. Minasyan, *Relatively hyperbolic groups are C^*-simple*, Journal of Functional Analysis, 243(1) (2007), 345-351.

[2] M. Bekka, M. Cowling, P. de la Harpe *Some groups whose reduced C^* algebra is simple*, Publications mathématiques de l'I.H.É.S., tome 80 (1994), p. 117-134.

[3] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, CA, 1994, 661 p., ISBN 0-12-185860-X.

[4] P. de la Harpe, *Groupes hyperboliques, algèbres dopérateurs et un théorème de Jolissaint*, C. R. Acad. Sci. Paris Sér. I Math. 307 (1988), no. 14, 771-774.

[5] P. de la Harpe. *On operator algebras, free groups, and other groups*, Astérisque 232 (S.M.F. 1995) 121-153. (1995), 121-153.

[6] S. Gong. *Finite part of operator K-theory for groups with rapid decay*, to appear in Journal of Noncommutative Geometry.

[7] P. Jolissaint. *Rapidly decreasing functions in reduced C^*-algebras of groups*, Trans. Amer. Math. Soc. 317 (1990), 167-196.

[8] P. Jolissaint. *K-theory of reduced C^*-algebras and rapidly decreasing functions on groups*, K-theory, (1989), no. 6, 723-735.

[9] P. Papasoglu. *Notes on hyperbolic and automatic groups*, `https://www.math.ucdavis.edu/~kapovich/280-2009/hyplectures_papasoglu.pdf`.

[10] R.T. Powers, *Simplicity of the C^*-algebra associated with the free group on two generators*, Duke Math. J. 42 (1975), 151-156.

[11] S. Weinberger, G. Yu *Finite part of operator K-theory for groups finitely embeddable into Hilbert space and the degree of non-rigidity of manifolds*, 2013.

[12] Z. Xie and G. Yu. *A relative higher index theorem, diffeomorphisms and positive scalar curvature*, Advances in Mathematics, 250 (2014) 35-73.

[13] Z. Xie and G. Yu. *Higher rho invariants and the moduli space of positive scalar curvature metrics*, arXiv:1310.1136, 2012